CAUTION

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Exteris Stressgard®

TURF FUNGICIDE

ACTIVE CONSTITUENTS: 12.5 g/L FLUOPYRAM

> 12.5 **TRIFLOXYSTROBIN** g/L

Distributed by CENTAUR

FUNGICIDE GROUP 11

5 L

For control of various diseases in turf as specified in the Directions for Use.

DIRECTIONS FOR USE RESTRAINTS

DO NOT apply by aircraft.

SPRAY DRIFT RESTRAINTS

DO NOT apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturer specifications that refer to ASAE S572 Standard to the BCPC Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are:

- 1. date and start and finish times of application;
- 2. location address and areas sprayed;
- **3.** full name of this product;
- 4. amount used per hectare and number of hectares applied to;
- 5. situation and disease;
- 6. wind speed and direction during application;
- 7. air temperature;
- 8. nozzle brand, model and type and spray system pressure measured during application;
- **9.** name and address of person applying this product.

(Additional record details may be required by the State or Territory where this product is used).

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas, including aquacultural ponds or surface streams and rivers, within 5 metres downwind from the application area.

CROP	DISEASE	RATE	CRITICAL COMMENTS
Turf	Fusarium or Microdochium patch (Microdochium nivale); Dollar spot (Sclerotinia homoeocarpa); Helminthosporium disease (Bipolaris spp., Drechslera spp., Exserohilum spp.); Brown patch (Rhizoctonia solani) Gray leaf spot (Pyricularia grisea) Rust (Puccinia spp.)	10 L/ha Water volume: 400-800 L/ha	Apply on a 14-28 day spray interval to established turf when conditions are favourable for disease development. Use shorter spray interval under high disease pressure or for early curative treatment. For Brown Patch control, apply sufficient irrigation to wash the product into the crown and upper root zone. Apply a maximum of four applications per year. Apply a maximum of 40 L / ha per year.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

DO NOT graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock.

GENERAL INSTRUCTIONS

Application

Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

IPM Compatibility

Risk to non-target insects: Exteris may have adverse effects on some non-target beneficial insects, in particular predatory bugs (e.g. *Orius* spp.). Safety to ladybirds and syrphid flies has not been evaluated. This risk should be taken into account where integrated pest management utilising these organisms is practised.

Compatibility

As formulations of other manufacturers' products are beyond the control of 2022 Environmental Science AU Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

RESISTANCE WARNING

GROUP 7 11 FUNGICIDE

Exteris Stressgard Turf Fungicide is a combination of a succinate dehydrogenase inhibitor (SDHI) and a quinone outside inhibitor (QoI) fungicide. For fungicide resistance management Exteris Stressgard is both a Group 7 and a Group 11 fungicide. Some naturally occurring individual fungi resistant to Exteris Stressgard and other Group 7 and Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Exteris Stressgard and other Group 7 and Group 11 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, 2022 Environmental Science AU Pty Ltd accepts no liability for any losses that may result from the failure of Exteris Stressgard to control resistant fungi.

PRECAUTIONS

Re-entry period Do not allow entry into treated areas until the spray has dried.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or water courses with this product or used containers.

PROTECTION OF LIVESTOCK

DO NOT graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS

May irritate the eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes. When handling product, wear chemical resistant gloves. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26)

SAFETY DATA SHEET

Additional Information is listed in the Safety Data Sheet which is available at www.au.envu.com.

CONDITIONS OF SUPPLY

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE RETURN AT ONCE UNOPENED.

To the extent permitted by law 2022 Environmental Science AU Pty Ltd does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. 2022 Environmental Science AU Pty Ltd does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter. To the extent only that 2022 Environmental Science AU Pty Ltd is prevented by law from contracting out of the Consumer Guarantees Act then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business

purposes.

APVMA Approval No.: 83704 / 108864

Envu, the Envu logo and Exteris Stessgard® are trademarks owned by Environmental Science U.S. LLC or one of its affiliates.

©2023 Environmental Science U.S. LLC

2022 Environmental Science AU Pty Ltd ABN 49 656 513 923 Suite 2.06, Level 2, 737 Burwood Road, Hawthorn East, Vic. 3123 Technical enquiries 1800 024 209 technicalsupport.australia@envu.com www.au.envu.com

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL +61 2 9037 2994 or 1800 862 115 toll free.

The toll free phone number is possibly accessible, but not guaranteed from payphones within Australia and is not accessible from outside of Australia.

Batch No. DOM:

drumMUSTER logo